

Behavioral Hearing Assessment in Babies & Infants

4th Phonak European Pediatric Amplification Conference November 14th, 2011



Anne Marie Tharpe

Vanderbilt University School of Medicine



Preview

- Background
- II. Behavioral Observation
- III. Visual Reinforcement Audiometry
- IV. Conditioned Play Audiometry



Background

In today's world with otoacoustic emissions, auditory brainstem responses, and auditory steady state responses, do we still need to learn behavioral test techniques?



- Auditory neuropathy/dysynchrony
- Fragile health care needs (30-40% of children have additional disabilities) many of whom cannot be sedated
- Physiologic equipment is expensive and may not be available



Age-Based Protocols

- Chronologically/Developmentally Birth to 4 months
- Chronologically/Developmentally
 5 24 months
- Chronologically/Developmentally
 25 60 months



ASHA Guidelines (2004)




II. Behavioral Observation






Wilson, Moore & Thompson, 1976





3-Month-Olds

Moore, Wilson, Thompson, 1977





III. Visual Reinforcement Audiometry





ASHA Protocol: 5-24 Months

- Speech & frequency-specific stimuli (.5, 1.0, 2.0 & 4.0 kHz)
- Bilateral testing with insert earphones





"It is easy to teach a baby the headturn. The hard part is teaching the baby <u>not</u> to turn."

(Nozza, 1999)



Control Trials


 An observational time when the examiner records whether a response occurs without a stimulus.


Should be the same amount of time allowed

for a response











Key Elements of a Conditioning Program:

- Appropriate stimulus
 - Waiting posture
 - Response
 - Reinforcer



Appropriate Stimulus

- Broad-band, complex
 - Frequency-specific



Waiting Posture








Response: 90° head turn





Reinforcer



(Moore, Wilson, Thompson, 1977)



MEDICAL CENTER





Suggested VRA Protocol:

✓ Starting level of 30 dB

✓ Step size of 20 dB down, 10 dB up

✓ No conditioning, unless indicated



IV. Conditioned Play Audiometry





ASHA Protocol: 25-60 Months

- Behavioral Assessment
 - Visual Reinforcement
 Audiometry (VRA)
 - Visually Reinforced
 Operant Conditioning
 Audiometry (VROCA)
 - Tangible Reinforcement
 Operant Conditioning
 Audiometry (TROCA)
 - Conditioned Play Audiometry (CPA)






Thompson, Thompson, & Vethivelu, 1989

- ✓ Compared effectiveness of 3 test procedures on 2-year-olds (VRA, VROCA, & CPA)
- √ 100% of the VRA group conditioned to task
- √ 83% of VROCA group conditioned to task
- √ 68% of CPA group conditioned to task



Responses of 2-yr-olds





Conclusions of Thompson et al.

- ✓ On average, more responses with CPA prior to habituation than with VRA or VROCA
- ✓ However, less likely to condition to CPA (~70%)
 than to VRA or VROCA



Thank you...



(anne.m.tharpe@vanderbilt.edu)